

Further additions to the knowledge of Indo-Pacific Mollusca in the Mediterranean Sea

(Lessepsian migrants)

By Al. Barash and Z. Danin

Summary ¹⁾

A report on 29 Indo-Pacific species found in the Mediterranean is presented. These species have not been discussed in the general reports on the migrations of Indo-Pacific species published by G. HAAS (1948), GHISOTTI (1974), BARASH and DANIN (1973) and in various articles on the migrating species collected during 1971–1984.

The 29 species recorded in the present report belong to 19 Gastropoda (Prosobranchia – 10, Opisthobranchia – 9) and to 10 Bivalvia (Pteriomorpha – 3, Heterodonta – 7).

The majority of the presented species (22) have been collected only close to the Suez Canal, along the Mediterranean coasts of Israel and Sinai. 7 species are recorded only or also from sites outside of Israel and Sinai: Black Sea, Aegean Sea, Adriatic Sea, Ionian Sea, Tyrrhenian Sea.

The greatest part of the Indo-Pacific species in the Mediterranean (79 of 91) are from the Erythrean subregion. A smaller part (12 species) were not recorded from the Erythrean subregion, but from areas in the Indo-Pacific remote from the Mediterranean. It is unlikely that these species would be able to reach the Mediterranean in larval stage, since the life span of molluscan larvae is rather limited. It is assumed that certain migrants were transported in adult stage by carriers (ships, fishes etc).

Of the 91 Indo-Pacific species appearing in the Mediterranean, 44 species have been collected as one or a few shells. The shells may be brought occasionally into the Mediterranean and it is doubtful if we can consider these species as inhabitants of this sea. 47 species were found live or in great numbers of shells, they may be regarded as residents in the Mediterranean.

The following 13 species may be qualified as permanent inhabitants in the Mediterranean: *Diodora rueppelli*, *Mimolia nedyma*, *Pirenella cailliaudi*, *Cerithium scabridum*, *Rhinoclavis kochi*, *Thais carinifera*, *Bursatella leachi savigniana*, *Hypselodoris infucata*, *Brachidontes variabilis*, *Pinctada radiata*, *Malleus regula*, *Paphia textile* and *Gastrochaena cymbium*.

Based on the data accumulated since the opening of the Suez Canal (1869) it may be stated: 1). The number of immigrating molluscan species has increased gradually during the past years. 2). The great majority of the Indo-Pacific immigrants are concentrated in the Eastern Mediterranean. A few penetrated the Western Mediterranean but did not surpass the Siculo Tunisian threshold.

Introduction

Several summary reports devoted to the migration of Indo-Pacific mollusca into the Mediterranean have been published during this century by G. HAAS (1948), GHISOTTI (1974 b) and BARASH & DANIN (1973, 1977). Numerous articles on separate Indo-Pacific species collected in the Mediterranean during 1971–1984 have been published by H. K. MIENIS and by other authors.

¹⁾ The species *Melibe fimbriata*, recorded in the addendum, was not included in the lists of the paper.

Today we are able to present an additional report on 29 Indo-Pacific species in the Mediterranean which have not been discussed in the general reports of 1948–1977 mentioned above. Eight species are recorded for the first time; 3 species were mentioned by name only (but not discussed) in the article by BARASH & DANIN (1982: 107) and 18 species have been dealt with in articles on individual Indo-Pacific immigrants.

Along with the 29 species not yet discussed in earlier reports, 4 species, which have been treated previously are included in this paper for supplementary data (see general remarks, p. 130ff.).

This report is based on material obtained from various sources, as follows:

A great deal of dredging was carried out during 1974–1977 by the late Prof. Ch. Lewinsohn (Tel Aviv University) and his assistants M. Tom and B. Galil. They worked in the infralittoral zone in north and south Israel and opposite the Mediterranean coast of north Sinai. Among the material collected by them the nudibranch *Plocamopherus ocellatus* is noteworthy because of its power of luminescence (O'DONOGHUE 1929: 808).

Investigations of the benthic fauna in the Haifa area, concerned with pollution, were carried out by Dr. Ch. Hornung of the Israel Oceanographic Limnological Research Institute. The molluscs collected there, mainly by diving, were put at the authors' disposal for examination. Amongst this material were the rare species *Cerithiopsis pulvis* and *Cingulina isseli* figured by SAVIGNY, 1817 (BOUCHET & DANRIGAL 1982).

Some Indo-Pacific species new to the Mediterranean have been revealed in the malacological collections of Israel such as *Planaxis punctostriatus*, *Vasum turbinellus*, *Ventomnestia girardi* and *Soletellina rubra* in the Zoological Museum of the Hebrew University, Jerusalem, *Divaricella angulifera* and *Trapezium oblongum* in the Department of Zoology, Tel Aviv University and *Cerithium nesioticum* in the collection of Dr. Yaron, Ben Gurion University, Beer Sheva.

Among the new species obtained from shore collectors, the most interesting was *Sphenia rueppelli*, which compares remarkably with the holotype of this species in the British Museum.

The specimens of the species enumerated in the present report are kept for the most part (20 species) in the collections of Tel Aviv University (NS) and the Hebrew University, Jerusalem (HJU). The remaining 9 species are deposited in the collections of the following scientific institutions abroad: Messina University, Civica Stazione Idrobiologica Milano; University of Ege, Izmir; British Museum, London, and in the private collections of Dr. J. J. van Aartsen (Holland), Mr. M. Bloecher (West Germany), Dr. F. Carrozza and Mr. Raybaudi Massilia (Italy).

The identification of 8 species reported for the first time in this paper was made by the following specialists: Gastropoda—Dr. J. J. van Aartsen, Dieren, Holland; Dr. F. Carrozza, Pisa, Italy; Dr. H. Gantes, University of Bordeaux, France; Dr. Mr. M. S. S. Lavaley, Rijksmuseum van Natuurlijke Historie, Leiden, Holland; Prof. T. E. Thompson, University of Bristol, England. Bivalvia—Prof. E. Fischer-Piette and Dr. P. Metivier of the Museum National d'Histoire Naturelle, Paris; Dr. R. N. Kilburn, Natal Museum, South Africa and Dr. Morris-Whybrow, British Museum, Natural History, London. The authors gratefully acknowledge their assistance.

Notes on the species

Remarks

1. The term "specimens" has been used for molluscs collected alive. Others are indicated as shells or valves.
2. Species marked by asterisk have been recorded in the earlier reports on Indo-Pacific species in the Mediterranean and the data on them are supplementary.

Gastropoda

Subclass Prosobranchia

Order Archaeogastropoda

Trochidae

Monodontinae

Clelandella infuscata (Gould, 1861)

DI NATALE 1982: 573, fig. 2

syn.: *Ziziphinus infuscatus* Gould, 1861; JOHNSON, R. J. 1964: 92

Cantbaridus infuscatus (Gould, 1861); KIRA 1968: 10, fig. 5

Mediterranean record (Sicily): DI NATALE 1982: 573

Material: One shell found in Ganzirri (Sicily, Straits of Messina), March 1972, 17 m depth, sand and small stones.

Distribution: Japanese Sea (Kira, 1968: 10).

Trochinae

Trochus (Infundibulops) erythraeus Brocchi, 1823*

THIELE 1931: 55

Mediterranean record: Israel – BARASH & DANIN 1973: 305

Material: 1. Live: Infralittoral zone, south of Netanya, 12. VII. 1971, 2–2.5 m depth, on rocks, 1 specimen (NS 19721); 12. XII. 1977, 2 m depth, 1 specimen juv. (NS 19722).

2. One shell: Shiqmona, south of Haifa, 2. XI. 1968, on the beach (NS 8897).

Remark: For the distribution of this species, see BARASH & DANIN 1973: 305.

Neritidae

Nerita sanguinolenta Menke, 1829

POR 1978: 102

syn. *Nerita forskalii* Recluz, 1841

Nerita crassilabrum Smith, 1885

Nerita albicilla Auct. (non Linne, 1758); MIENIS 1974: 45

Neritina kinzelbachi Nordsieck, 1973

Mediterranean record (Greece); Nordsieck 1973: 4

Material: One shell at Karpathos Is.

Distribution: Suez Canal (MOAZZO 1939: 200); Gulf of Suez and Aqaba (LAMY 1938: 78). Red Sea, Perim Is. (STURANY 1903: 56). Aden (SMITH 1891: 419).

Remark: NORDSIECK (1973: 4) described one shell found at Karpathos Is. as a new species under the name *Neritina kinzelbachi* spec. nov. According to MIENIS (1974: 45) the figure of *N. kinzelbachi* is identical with *Nerita sanguinolenta* (Menke, 1829), a common erythraean species.

Order Mesogastropoda

Planaxidae

Planaxis punctostriatus Smith, 1872

MIENIS 1981: 361

syn. *Planaxis lineolatus* Gould, 1849 (not *Planaxis lineolatus* Risso, 1826); LAMY 1938: 70; MIENIS 1980a

Mediterranean record (Israel): MIENIS 1981: 361

Material: South of Dor, 1950, 3 shells, in the collection of Prof. G. Haas, Hebrew University, Jerusalem (HUJ 21, 385).

Distribution: Gulf of Aqaba (SAFRIEL & LIPKIN 1964: 189); Gulf of Suez (LAMY 1938: 70); Madagascar (DAUTZENBERG 1929: 493); Siam (P. H. FISCHER 1970: 101); Solomon Is. (ABBOTT 1958: 204). Tuamotu and Manihiki Is. (JOHNSON 1964: 102).

Cerithiidae

Cerithium nesioticum Pilsbry and Vanatta, 1906

CERNOHORSKY 1972: 69, Plate 16, fig. 1

syn. *Semivertagus nesioticus* (Pilsbry & Vanatta, 1906); HABE 1970: 43, plate 13, fig. 2

Cerithium lacteum Kiener, 1841; TRYON 1887: 143, plate 27, figs. 29, 30, 33

Liocerithium lacteum (Kiener, 1841); SOLEM 1953: 220

Mediterranean record (Israel): MIENIS 1977 a: 45

Material: Shiqmona (south of Haifa), January 1971, on the beach, one shell (MIENIS, 1977 a: 45)

Distribution: Gulf of Suez and Aqaba (LAMY 1938: 66), Red Sea (STURANY 1903: 53, 70). Gulf of Aden, Perim, Djibouti, Obock (JOUSSEAUME 1930: 284). Madagascar (DAUTZENBERG 1929: 274 [480]). Mauritius (VIADER 1937: 42). East Africa – Zanzibar, Ceylon, Cocos Keeling Is. (ORR MAES 1967: 114). Indonesia (ADAM & LELOUP 1938: 106). Philippines (MCANDREW 1870: 441). Japan (HABE 1970: 43). Central Pacific Islands: New Caledonia, Palau Islands, Cook Islands (ORR MAES 1967: 114). Tahiti, Tuamotu Islands (DAUTZENBERG & BOUGE 1933: 308), Hawaii Islands (KAY 1979: 123).

Remark: The name *Cerithium lacteum* Kiener, 1841, has priority but was preoccupied by *Cerithium lacteum* Philippi, 1836.

Cerithiopsidae

Cerithiopsis pulvis (Issel, 1869)

PALLARY 1926: 66, plate 4, fig. 5

syn. *Cerithium* (*Cerithiopsis*?) *pulvis* Issel, 1869: 150; BOUCHET & DANRIGAL 1982: 14, fig. 35

Mediterranean record (Israel): AARTSEN & CARROZZA 1983: 37

Material: Haifa, VII. 1980, 9 m depth, 2 shells and one fragment, 10. V. 1982, 12 m depth, 3 shells; Hadera, VII. 1982, 9 m depth, 3 fragments. All the material was obtained by diving. The largest specimen measures 4 × 1 mm and has 7.5 teleconch whorls (AARTSEN & CARROZZA 1983: 40).

Distribution: Gulf of Suez (ISSEL, 1869: 151).

Hipponicidae

Hipponix (Sabia) conicus (Schumacher, 1817)

WENZ, 1960–1962 (1938–1944): 888, fig. 2612

syn. *Amalthea conica* Schumacher, 1817; KURODA, HABE & OYAMA 1971: 84

Sabia conica (Schumacher, 1817); KAY 1979: 179

Hipponix australis (Quoy and Gaimard, 1834); STURANY 1903: 49 (257); KURODA and al. 1971: 84

Record: New to the Mediterranean (Israel) (Det. Kilburn)

Material: Dor, 30. IX. 1980, by diving, 1–2 m depth, one shell with the rest of soft body (NS 19723).

Distribution: Gulf of Suez (MOAZZO 1939: 191); Gulf of Aqaba (LAMY 1938: 69). Red Sea (STURANY 1903: 49 (257); Aden (SMITH 1891: 418); South Africa-Natal (KILBURN & RIPPEY 1982: 56); Madagascar (DAUTZENBERG 1929: 300); Mauritius (VIADER 1937: 48); Indonesia (ADAM & LELOUP 1938: 108); Japan (KIRA 1968: 28); Australia (ALLAN 1950: 85); Central Pacific Is. (DAUTZENBERG & BOUGE 1933: 371); Hawaii Is. (KAY 1979: 179); East Pacific; British Columbia (COWAN 1974: 3).

Remark: The small limpet *Hipponix conicus* (Schumacher) is a very common Indo-Pacific species that lives epizootically on the shells of larger gastropods. Methods of feeding are described in YONGE 1953: 11; TINKER 1967: 44; CERNOHORSKY 1968: 275 and MASTALLER 1979: 58.

Strombidae

Strombus decorus (Roeding, 1798)

ABBOTT R. T., 1960: 137, 139, plate 14, figs. 13, 14

Mediterranean record (Turkey): L. R. MASSILIA, 1983: 20; M. BLOECHER (in Nicolay, K. & E. R. MANOJA) 1983: 17, 18

Israel: H. K. Mienis (pers. comm.)

Material: I. Turkey – 1) Bozborum, summer 1983, one shell and 2 live specimens between Gelibolu and Marmaris, 12 specimens, (Leg: L. R. Massilia)

2) Between Silifke and Mersin, autumn 1983, many specimens (Leg. M. Bloecher)

II. Israel – Shiqmona (south of Haifa), one shell in 1983, 3 shells, February-March, 1984 (Leg. B. Singer).

Distribution: 1) *Strombus decorus decorus* (Roeding, 1798) – Zanzibar, Mozambique, South Africa, Saudi Arabia, Madagascar, Indian Ocean Islands, Ceylon, Thailand, Indonesia, Burma (ABBOTT, T. 1960: 137, 138)

2) *Strombus decorus persicus* Swainson, 1821 – Persian Gulf (ABBOTT, T. 1960: 139); Gulf of Aden (NICOLAY & MANOJA 1983: 18)

3) *Strombus decorus raybaudi* n. ssp. – Turkey (Mediterranean) (NICOLAY & MANOJA 1983: 18).

Remarks: The specimens of *Strombus decorus* collected in Turkey and Israel have great similarity to *S. decorus persicus*

Some differences from *S. decorus persicus* are evident in the specimens of *Strombus* collected in Turkey, according to NICOLAY & MANOJA (1983). In their opinion a taxonomic separation from *S. decorus persicus* is necessary, and they established a new subspecies *S. decorus raybaudi*.

The existence of *Strombus* species in the Mediterranean was previously considered as doubtful. Only *Strombus lentiginosus* was recorded (AHARONI 1934: 476) but no information was available about the number of specimens collected, nor of the place of their deposition.

Cypraeidae

Erosaria turdus (Lamarck, 1810)

TAYLOR & WALLS 1975: 263

syn. *Cypraea turdus* Lamarck, 1810; PALLARY 1926: 96, fig. 31;

BURGESS 1970: 235, plate 22, fig. 8

Record: New to the Mediterranean (Israel) – (Det. Burgess)

Material: Dor, 6. VI. 1980, in shallow water, 1–1.5 m depth, one shell juv., length 37 mm, width 24 mm, height 19 mm.

Distribution: Suez Canal – Little Bitter Lake (MOAZZO 1939: 166, 272); Gulfs of Suez and Aqaba, Red Sea (STURANY 1903: 47); Aden (SHOPLAND 1902: 175); Kenya (BURGESS 1970: 235); Madagascar (DAUTZENBERG 1929: 259); Arabian Sea – Karachi (MELVILL & STANDEN 1901: 384); Gulf of Oman (BURGESS 1970: 235); Persian Gulf (BIGGS 1973: 365); Ceylon, Indonesia (JICKELI 1884: 197).

Palmadusta gracilis (Gascoïn, 1849)

TAYLOR & WELLS 1975: 262

syn. *Cypraea gracilis* Gascoïn, 1849; BURGESS 1970: 140, plate 9a, fig. A

Mediterranean records: 1) Turkey – BLOECHER 1983: 26

2) Israel – MIENIS & SINGER 1983: 535

Material: 1) Mediterranean coast of east Anatolia (Turkey), between the small towns of Silifke and Mersin, rocky shore.

Live: Infralittoral, 3 m depth, autumn 1983, under stone, one specimen, shell measures: 16.4×10×7.9 mm (BLOECHER 1984).

One shell, near the shore, on a rock, 28. X. 1982 (leg. BLOECHER).

2) Mediterranean coast of Israel

Live: Hadera, 25. VII. 1981, 1 m depth, one specimen (NS 19724).

Shells: On beaches – Akhziv, 28. VII. 1983, 4 shells, Nahariyya, 28. VII. 1983, one shell, Akko, VIII, 1983, one shell, Netanya, 8. XI. 1983, one shell, Palmahim, 3. VII. 1983, 15. VII. 1983, 5. VIII. 1983, 3 shells.

Distribution: Gulf of Suez (STURANY 1903: 68), Gulf of Aqaba (MIENIS 1971: 30); Red Sea (MASTALLER 1978: 132); Kenya, Oman (BURGESS 1970: 140); Persian Gulf (SMYTHE 1979: 67); South India-Gulf of Manaar (SATYA-MURTI 1952: 127); China, Japan (BURGESS 1970: 140); Indonesia-Java (REGTERN 1945: 146); Australia (CATE 1968: 291); Fiji Is. (CERNOHORSKY 1967: 95); Hawaii Is. (KAY 1979: 202).

Remarks: *Palmadusta gracilis notata* (GILL 1858) is the common subspecies in the Erythraean area – (MIENIS 1971: 29). In some publications *P. gracilis* has been reported as *Palmadusta fimbriata* (Gmelin) e. g. by MCANDREW 1870: 440, STURANY 1903: 276 and MOAZZO 1939: 164.

The recent findings of *P. gracilis* in 1982 and 1983 in two areas of the Mediterranean (Israel and South Turkey) are quite remarkable. The possibility of permanent settlement and colonisation of this Indo-Pacific species in the Mediterranean is most probable.

Order Neogastropoda

Muricidae

Rapana venosa (Valenciennes, 1846)

GHISOTTI 1974b: 12, fig. 38

syn. *Purpura venosa* Valenciennes, 1846

Rapana bezoar Reeve, 1847 (non Linnaeus, 1767), GHISOTTI 1974a: 125

Rapana thomasiana Crosse, 1861

Rapana pontica Nordsieck 1969; MIENIS 1976a: 39

Records: *Rapana venosa* is an introduction from the Far East, which reached the Black Sea before 1947. Since then it has also been reported from the Adriatic Sea (MIENIS 1976a: 41) and recently from the Tyrrhenian Sea – Elba Is. (TERRENI 1980: 11).

Material: Data on the collecting of *Rapana venosa* are presented in the following publications: in the Black Sea (GROSSU & LUPU 1964); in the Adriatic Sea (GHISOTTI 1971a, 1974a; MEL 1976) and in the Tyrrhenian-Elba Is. (TERRENI 1980). The live specimens collected in the areas indicated were obtained in the infralittoral zone, at depths of 12–90 m. The measurements of the specimens are various; generally large. The dimensions of a specimen collected from the Elba Is. are: height – 89 mm, width – 68 mm, and of a specimen taken from the Gulf of Trieste, height – 110 mm, width – 75 mm.

Distribution: Japan-Tartari Canal (GHISOTTI 1971a: 55), Yellow Sea and Pacific Ocean (GROSSU & LUPU 1964: 215).

Remarks: *Rapana venosa* obviously covered the long route from the Far East to the Mediterranean through human intervention (GHISOTTI 1974a: 125). The occurrence of *R. venosa* in the Black Sea was confirmed again recently by Prof. Grossu and in the Adriatic Sea by Dr. Ghisotti (in litt.).

Rapana venosa seems already to be past the phase of acclimatization in the Mediterranean and is able to reproduce in the new environment (GROSSU & LUPU 1964: 216).

Vasidae

Vasum turbinellus (Linnaeus, 1758)

CERNOHORSKY 1972: 164, plate 49, figs. 2, 2a

Mediterranean record (North Sinai): MIENIS 1973: 6

Material: North Sinai, near Bardawil Lagoon, XII. 1956, one shell.

Distribution: Gulf of Aqaba (LAMY 1938: 49); Gulf of Suez, Red Sea (STURANY 1903: 39); Aden (SHOPLAND 1902: 173); Obock (JOSSEAUME 1888: 182); East Africa-Dar es Salaam (SPRY 1961: 25); Madagascar (DAUTZENBERG 1929: 196); Ceylon (ROBERTSON 1969: 5); Cocos Keeling Is. (ABBOT 1950: 84); Indonesia, New Guinea (ADAM & LELOUP 1939: 114); Philippines, Marianos Is. (HIDALGO 1905: 6); Japan (KIRA 1968: 84); Solomon Is. (SOLEM 1953: 225); Australia (WILSON & GILLETT 1971: 112; RIPPINGALE & MCMICHAEL 1961: 126).

Subclass Opisthobranchia

Order Pyramidellomorpha

Pyramidellidae

Chrysallida fischeri (Hornung and Mermod, 1925)

AARTSEN & CARROZZA 1979: 29, figs. 1A, 1B

syn. *Pyrgulina fischeri* HORNUNG & MERMOD 1925: 27, fig. 1A

Mediterranean record (Israel): Aartsen & Carrozza 1979: 29

Material: Haifa Bay, 24. XII. 1974, 40–70 m depth, mud, several shells, average length 2.60 mm, width 0.80 mm (specimens kept in the collections of Aartsen & Carrozza).

Distribution: Gulf of Aqaba-Elat (Gittenberger by litt.); Red Sea-Massawa (Hornung & Mermod 1925: 27). This species has not been recorded from the Suez Canal.

Remark: In addition to *Chrysallida fischeri*, four other Indo-Pacific species of Pyramidellidae have been found in the Mediterranean: *Chrysallida maiae*, *Kleinella fulva*, ? *Monoptygma sinuata* and *Cingulina isseli*. The hosts of these ectoparasitic snails which feed upon invertebrates (as do Pyramidellidae in general) are as yet unknown.

Kleinella fulva (A. Adams, 1851)

LAVALEYE & BARASH 1981: 87, fig. 1

syn. *Monoptygma fulva* A. Adams, 1851

Pyramidella (Actaeopyramis) fulva (A. Adams) LAVALEYE & BARASH 1981: 87

Mediterranean record: (Israel). LAVALEYE & BARASH 1981: 87

Material:

Live: Haifa Bay, 14. XI. 1982, 9 m depth, one specimen, measuring height – 11.5 mm, width – 3.1 mm, height of last whorl – 5.1 mm, height of aperture – 3.1 mm. (NS 19725).

Shell: Atlit-Dor, south of Haifa, 15. II. 1967, 22 m depth, one shell (SFRS¹ 1357), measuring, height – 8.9 mm, width – 2.7 mm, height of last whorl – 4.8 mm, height of aperture – 2.9 mm.

Distribution: Arabian Sea: Karachi, Bombay (MELVILL & STANDEN 1901: 392); Japan, Philippines (LAVALEYE & BARASH 1981: 88).

¹) Sea Fisheries Research Station, Haifa

? *Monoptygma sinuata* Gould, 1861

JOHNSON, 1964: 150

A record new to the Mediterranean (Israel); determined tentatively by van Aartsen
Material:

Live: Haifa Bay, 14. XI. 1982, 9 m depth, 4 specimens, average measurements, height – 6.4 mm, width – 2 mm, height of last whorl – 4 mm, height of aperture – 3 mm (NS 19726).

Shell: Haifa Bay, 16. II. 1975, 73 m depth, mud, one shell.

Distribution: China Seas (JOHNSON 1964: 150).

Remark: “This species belongs to Actaeopyramis/Kleinella group of Pyramidellidae. The only Actaeopyramis which looks reasonably close seems to be *Monoptygma sinuata* Gould, 1861. In my opinion it is not one of the better known species of Actaeopyramis: *fulva*, *amoena*, *casta*, *lauta* or *eximia*. I have not been successful in a certain identification up to now” (AARTSEN, in litt. 4. XI. 1982).

Cingulina isseli (Tryon, 1886)

AARTSEN & CARROZZA 1983: 38, fig. 1

syn. *Eulimella cingulata* Issel, 1869; ISSEL 1869: 182, 332; BOUCHET & DANRIGAL 1982: 12, fig. 73

Odetta cingulata Issel; PALLARY 1926: 58, plate 3, fig. 25

Mediterranean record: (Israel). AARTSEN & CARROZZA 1983: 38

Material: Haifa Bay, VII. 1980, 9 m depth, 8 shells; 10. V. 1982, 12 m depth, one shell. Measurements of largest shell collected are, height – 3 mm, width – 1 mm.

Distribution: Gulf of Suez (Issel 1969: 182); Red Sea-Massawa (AARTSEN & CARROZZA 1983: 38).

Order Bullomorpha (Cephalaspidea)

Bullidae

Bulla ampulla Linnaeus, 1758

HABE 1970: 136, plate 42, fig. 17

syn. *Bullaria ampulla* (Linnaeus, 1758); THIELE 1931: 384, fig. 483

Mediterranean record (Israel): BARASH & DANIN 1982: 107

Material: Gaash (north of Tel Aviv), 20. IV. 1978, on the beach, one shell (NS 19727).

Distribution: Gulf of Suez and Aqaba, Red Sea (STURANY 1903: 60, 74); Aden, Obock (JOUSSEAU 1888: 170); East Africa-Dar es Salaam (SPRY 1961: 29); Madagascar (DAUTZENBERG 1929: 365); Mauritius (VIADER 1937: 3); Persian Gulf (MELVILL & STANDEN 1901: 456); East Indies-Gulf of Manaar (SATYAMURTI 1952: 218); Ceylon (STARMUEHLNER 1974: 63); Cocos Keeling Is. (ABBOTT 1950: 86); Indonesia (ADAM & LELOUP 1938: 198); Philippines (HIDALGO 1905: 14); Japan (HABE 1970: 136); New Guinea (Papua Is) (TAPPARONE CANEFRI, 1874: 14); Solomon Is. (SOLEM 1953: 217); Gambier (DAUTZENBERG & BOUGE 1933: 53).

Remark: *Bulla ampulla* has been found live in the waters of Alexandria, according to EHRENBERG (1831). ISSEL (1869) remarked with regard to this record: “We hardly believe that it has been collected there, since it was never encountered in the Mediterranean by any other conchologist”.

Retusidae

Ventommestia girardi (Audouin, 1827)

MIENIS 1976 b, fig. 1

syn. *Cylichna girardi* (Audouin, 1827); PALLARY 1926: 74, plate 5, fig. 3; MOAZZO 1939: 135, fig. 10

Bulla girardi Audouin, 1827; BOUCHET & DANRIGAL 1982: 13, fig. 52

Alicula girardi (Audouin); ISSEL 1869: 341

Mediterranean record (Israel): MIENIS 1976b: 209

Material: Bardawil Lagoon (North Sinai), 9. I. 1974, on the beach, 3 shells, height – 3.0–3.2 mm.

Distribution: Suez Canal, Gulf of Suez (MOAZZO 1939: 135); Gulf of Aqaba (LAMY 1938: 86); Red Sea, China Sea, Ryu Kyu Is. (Japan) (MIENIS 1976b: 210).

Order Aplysiomorpha (Anaspidea)

Aplysiidae

Bursatella leachi de Blainville, 1817; *subspec. savigniana* Audouin, 1826*

BARASH & DANIN, 1973: 329

Material:

Israel. Haifa Bay, 10. IV. 1979, 80 m depth, mud, 1 specimen (NS 16597); 22. VI. 1981, sea water pond (at Electricity Station, Haifa), 1–2 m depth, 1 specimen (NS 19010); Maayan Zvi, 28. VI. 1981, sea water pond, 1–2 m depth, 3 specimens juv., (NS 19011); Maagan Michael, 31. I. 1978, sea water pond, 1–2 m depth, 1 specimen (NS 16570); Off Bardawil, 1. IV. 1978, 14.5 m depth, 1 specimen juv. (NS 16398).

Greece: Salamis Is. – east coast, VIII–X. 1977, many specimens, in shallow water, among algae (*Ulva*) (Dr. Alice Panou, in litt.); Khios Is. 16. IV. 1975, 36 m depth, silt, collection of Thessaloniki University (75–G₂748). Lesbos Is.-Kalloni, 3 m depth, seagrass meadow (? *Posidonia*), collection of Thessaloniki University (116–G₂748).

Italy: Sicily–Syracuse, 37° N, Bay of Panagia, 10. VII. 1978, 22 m depth, mud, 1 specimen, 8 cm long (in the collection of the Hydrogeological Station of Milano) (cat. no. 1547M) (PIANI 1980: 7). Gulf of Palermo, 38° N, 4–5 m depth, I. 1976, 1 specimen, III. 1976, 6 specimens, average length 7–13 cm, (Catalano, PARINELLO & DE LEO 1978: 123, 124). Gulf of Taranto, salt lakes Mar Grande and Mar Piccolo, 1970–1973, 4–9 m depth, sandy detritic bottom, often among algae (*Gracilaria*), 49 specimens, average length 3–9 cm. Egg strings deposited on *Gracilaria*, March to May (TORTORICI & PANETTA 1977: 256 and pers. comm. with P. Panetta).

Remark: The sea water ponds along the shores of Israel seem to be favourable habitats for *Bursatella leachi savigniana*. Adults and juveniles are often found there and the deposition of egg strings has been observed. Some ecological data for these ponds are: depth 1–2 m, bottom – sand mixed with silt, yearly temperature 12–35°C, 4°C difference between day and night, salinity 30–38‰, pH 7.6–8.4.

Order Nudibranchia

Family Triophidae (Kaloplocamidae)

Plocamopherus ocellatus Rueppell and Leuckart, 1828

O'DONOGHUE, 1929: 735, 806

syn. *Plocamophorus ocellatus* F. S. Leuckart, 1828; CHENU 1859: 403, fig. 3043; THIELE 1931: 426

Mediterranean record: (Israel); BARASH & DANIN 1982: 107

Material: Nizanim, 12. IV. 1977, 50 m depth, 1 specimen (NS 19728)

Distribution: Suez Canal–Great Bitter Lake, Canal proper (O'DONOGHUE 1929: 806); Gulf of Suez (ISSEL 1869: 159); Red Sea (PRUVOT-FOL 1933: 100).

Chromodorididae

Hypselodoris infucata (Rueppell and Leuckart, 1828)*

BERTSCH & JOHNSON 1981: 62, 63, 2 figs.

syn. *Doris infucata* Rueppell and Leuckart, 1828; ISSEL 1869: 156

Glossodoris infucata (Rueppell and Leuckart, 1828); O'DONOGHUE 1929: 724

Glossodoris runcinata (Bergh, 1877); ENGEL & EEKEN 1962: 24
Chromodoris runcinata Bergh, 1877; RUDMAN 1971: 386

Mediterranean records (Israel): 1) Under the name *Glossodoris runcinata*: BARASH & DANIN 1974: 98; 2) Under the name *Hypselodoris infucata*: MIENIS & GAT 1981: 51

Material: Shiqmona, VI. 1983, 2–3 m depth, 1 specimen; Dor, 10. IX. 1974, 1 m depth, 1 specimen (NS 12914); 13. VIII. 1980, 1 m depth, 1 specimen (NS 981); Hadera, 25. I. 1971, 35 m depth, 1 specimen (NS 18944); Mikhmoret, 19. VI. 1978, 1 specimen; Ashdod, 8. II. 1968, 1 specimen (HUJ SLM 495); Ashkelon, 12 m depth, rocky ground, 7. IV. 1977, 1 specimen; 19. VI. 1978, 1 specimen; 20. VI. 1980, 3 specimens; 20. IX. 1980, 1 specimen; 27. IX. 1980, 1 specimen; 15. III. 1981, 1 specimen; 3. IV. 1981, 5 specimens; 11. IV. 1981, 15 specimens; 20. IV. 1981, 10 specimens; 7. V. 1981, 15 specimens (Data on Ashkelon given by G. Gat in litt.).

Distribution: Gulf of Suez-Tor (ISSEL 1869: 156); Gulf of Aqaba-Elat (ENGEL & EEKEN 1962: 24); Red Sea, East Africa-Zanzibar, Dar-es-Salaam, South Africa, Philippines Is., South Australia, New Caledonia, Fiji, Hawaii Is. (RUDMAN 1974: 387; BERTSCH & JOHNSON 1979: 42).

Remark: The numerous records of *H. infucata* from different localities along the Mediterranean coast of Israel and Sinai prove that this species has established itself permanently in this part of the eastern Mediterranean.

According to BERTSCH & JOHNSON (1981: 62) *H. infucata* feeds on the sponge *Dysidea fragilis*. In Ashkelon many specimens of *H. infucata* were observed on a bed of the sponge *Clathrina coreacea* (GAT & FAINZILBER 1983: 17).

Dendrodorididae

Dendrodoris nigra Stimpson, 1855

THOMPSON 1975: 504, fig. 5f, g

syn. *Doris nigra* Stimpson, 1855; O'DONOGHUE 1929: 730, 827

A record new to the Mediterranean (Det. J. Orr)

Material: North of Dor, 23. VII. 1980, 5 m depth, under stone, 1 specimen (NS 18713)

Distribution: Suez Canal-Little Bitter Lake, Kabret (O'DONOGHUE 1929: 827); Red Sea (PRUVOT-FOL 1933: 100); Madagascar (MARCUS & MARCUS 1970: 206); Persian Gulf (THORSON 1940: 221); India, Maldives Is. (SATYAMURTI 1952: 243); Hong Kong (ORR 1980: 112); Vietnam, Japan, Australia (THOMPSON 1975: 504); Central Pacific Is., Gilbert Is., New Caledonia (MARCUS & MARCUS 1970: 206); Hawaii Is., (KAY 1979: 473). *Dendrodoris nigra* is widely distributed in the Indian and Pacific Oceans.

Bivalvia

Subclass Pteriomorpha

Order Arcoida

Arcidae

Arcinae

Acar plicatum (Dillwyn, 1817)

KURODA, HABA & OYAMA 1971: 330, plate 67, fig. 17

syn. *Arca plicata* (Chemnitz) Dillwyn, 1817; BARNARD 1964: 372, fig. 2a; SMYTHE 1972: 495

Barbatia plicata (Dillwyn); DIGERONIMO & ROBBA 1979: 177, plate 1, fig. 2

A record new to the Mediterranean (Israel) – (Det. S. Morris Whybrow)

Material: Netanya, 14. II. 1978, intertidal zone, under stone, 1 specimen with rest of soft body (NS 19729)

Distribution: Suez Canal-Bitter Lake, Port Taufiq (MOAZZO 1939: 55, 257); Gulf of Suez and Aqaba (STURANY 1899: 35, 41); Aden (SHOPLAND 1902: 178); Djibouti, Obok (LAMY 1917: 7); Somalia (DIGERONIMO & ROBBA 1979: 177); South Africa, Madagascar (BARNARD 1964: 372); Seychelles, Oman (LAMY 1907: 87); Persian Gulf (SMYTHE 1972: 495); Japan (KIRA 1968: 121); Philippines, Indonesia, New Caledonia, Australia, New Zealand (PRASHAD 1932: 51, 52), Hawaii Is. (KAY 1979: 501).

Anadarinae

Scapharca demiri Piani, 1981

PIANI 1981: 284

syn. *Arca amygdalum* Philippi, 1847; DEMIR 1977: 197 figs. 1, 9

Mediterranean record (Turkey): DEMIR 1977: 197

Material: Izmir Harbor, in the spring, 1978, 5–15 m depth, sandy mud, mud, numerous specimens collected by Dr. A. Kocatas, University of Ege, Izmir.

Distribution: China Seas (LAMY 1907: 235). This species has not been recorded from the Suez Canal.

Remarks: According to Dr. B. Metivier of the Museum National d'Histoire Naturelle, the species common in the Izmir harbour is *Arca amygdalum* Philippi (DEMIR 1977: 198)

A new name *Scapharca demiri* was proposed for *Arca amygdalum* Philippi, since Link 1807 described other species under the same name *Arca amygdalum* (PIANI 1981: 284).

Scapharca inaequivalvis (Bruguere, 1789)

KURODA, HABE, OYAMA 1971: 333, plate 69, figs. 3, 4

syn. *Arca inaequivalvis* Bruguere, 1789; LAMY 1907: 256

Mediterranean record: (Adriatic Sea): GHISOTTI 1974: 11, under the name of *Scapharca cf. cornea* (Reeve)

Material: A few small specimens of *Scapharca inaequivalvis* appeared for the first time in 1969 and 1971 in the littoral zone of Ravenna (northern Adriatic) (GHISOTTI 1972: 21). Through the years the number of specimens found in the Adriatic Sea multiplied and became distributed southwards. Specimens have been found in Ancona and in 1981 reached Manfredonia (S. Adriatic). The range of vertical distribution of specimens collected is from shallow water near the shore up to a depth of 10 m. The dimensions of the specimens of different ages are: from 1 mm length and 0.6 mm height, up to adults of 73.5 mm length and 61 mm height (RINALDI 1977: 199–201).

Distribution: Philippines (LAMY, 1907: 256); Japan (KURODA, HABE, OYAMA, 1971: 334).

Remark: This species was at first recorded under the name *Scapharca cornea* Reeve, 1844 (GHISOTTI, 1973: 68). GHISOTTI & RINALDI (1976: 13) later concluded that the correct name of this new Indo-Pacific species should be *Scapharca inaequivalvis* (BRUGUIERE, 1789).

Subclass Heterodonta

Order Veneroida

Lucinidae

Divaricella angulifera (von Martens, 1880)

DAUTZENBERG, 1929: 418; LAMY 1938: 31

syn. *Lucina angulifera* von Martens, 1880; LAMY 1916a: 15

? *Divaricella quadrisulcata* (D'ORBIGNY 1846); LAMY 1916a: 14; MOAZZO 1939: 114

Mediterranean record (Israel): MIENIS 1979: 200

Material: Netanya, 9. IV. 1976, 1 m depth, 1 valve, length and height 12.5 mm (NS 19730); off Bardawil, 7. XII. 1956, depth unknown, 1 valve, length and height 12 mm (NS 19731).

Distribution: Gulf of Suez (LAMY 1938: 31); Gulf of Aqaba and Red Sea (STURANY 1899: 285); Djibouti (MASTALLER 1979: 148); Madagascar (DAUTZENBERG 1929: 418 (624); Mauritius and Seychelles (BARNARD 1964: 479); Indonesia (PRASHAD 1932: 160).

Remark: There has been a great deal of confusion regarding the proper name of this species. VON MARTENS 1880 described under the name *Lucina angulifera* specimens collected in the Red Sea, Seychelles and Mauritius (LAMY 1920: 272). Some other authors (STURANY 1899; MOAZZO 1939) assigned specimens collected in the Red Sea to *Lucina quadrisulcata* d'Orbigny known to occur in the western Atlantic. Priority therefore goes to the latter species. However, *Divaricella quadrisulcata* is different from *D. angulifera* "the divaricate impressed lines are less angulated" (MIENIS 1979: 200) and these two species are not synonymous.

Cardiidae

Laevicardium flavum (Linnaeus, 1758)

FISCHER-PIETTE 1977: 29

syn. *Cardium flavum* Linnaeus 1758; SATYAMURTI 1956: 91, plate XIV, figs. a, c

Vasticardium flavum (Linnaeus); KIRA 1968: 156, plate 56, fig. 7

Trachycardium flavum (Linnaeus); SPRY 1964: 29, plate II

Cardium pectiniforme Born, 1780; PRASHAD 1932: 266

Cardium rugosum Lamarck, 1819; FISCHER-PIETTE 1977: 30

Cardium orbita Broderip & Sowerby, 1833; DAUTZENBERG & BOUGE 1933: 446

Trachycardium hawaiiensis Dall, Bartsch & Rehder, 1938; KAY 1979: 556

A record new to the Mediterranean – (Det. Fischer-Piette)

Material: Dor, 6. VI. 1980, intertidal zone, 1 valve, length 40 mm, height 48 mm (NS 19732)

Distribution: Gulfs of Suez and Aqaba, Red Sea proper (STURANY 1899: 39), Aden (SHOPLAND 1902: 178); East Africa-Dar es Salaam (SPRY 1964: 29); Seychelles (FISCHER-PIETTE 1977: 32); Madagascar (DAUTZENBERG 1929: 382 [588]); Mauritius (VIADER 1937: 68); Gulf of Oman, Persian Gulf (MELVILL & STANDEN 1906: 837); India, (Manaar) (STARBUHLNER 1974: 68); Indonesia (PRASHAD 1932: 266); Philippines, Jolo Archipelago and Marianas Is. (HIDALGO 1905: 15); Japan (KIRA 1968: 156); Tuamotu (LAMY 1906: 214); Australia, north and south (RIPPINGALE & MCMICHAEL 1961: 186); New Guinea (FISCHER-PIETTE 1977: 32); Hawaii Is., (KAY 1979: 556).

Remark: *Laevicardium flavum* is a wide ranging species in the Indo-West Pacific. It is extremely variable and thus also appears under numerous synonyms.

Psammobiidae

Soletellina rubra (Schroeter, 1788)

MIENIS 1980b: 301

syn. *Solen ruber* Chemnitz, 1782; MASTALLER 1979: 165

Soletellina rueppelliana (Reeve, 1857); MIENIS 1980b: 301

Psammotella rueppelliana Reeve, 1857; LAMY 1918: 244

Solen roseus Gmelin, 1791

Soletellina rosea (Gmelin, 1791); STURANY 1899: 277

Mediterranean record (Israel): MIENIS 1980b: 301

Material: Off Bardawil, VII. 1967, by dredging, depth unknown, 1 shell, length 54 mm, height 26 mm, 1 damaged valve (NS 19734); on the beach at Bardawil Lagoon, 7. IV. 1968, 2 shells, length 54.4 mm, height 27.4 mm (MIENIS 1980b: 301)

Distribution: Suez Canal: Lake Timsah, Great and Little Bitter Lakes, Canal proper (MOAZZO 1939: 97, 264); Gulf of Suez, Red Sea (STURANY 1899: 38 [292]); Aden, Djibouti (LAMY 1918: 245); Kenya, Persian Gulf (fide Fischer-Piette, by litt.).

Remark: This species from the Red Sea is today acclimatised in Port Said (PALLARY 1912: 175, plate XVII, figs. 3, 4).

Soletellina subradiata (Reeve, 1857)

Fischer-Piette (unpublished manuscript on Psammobiidae)

syn. *Psammotella subradiata* (Desh. m. s.) in Reeve, 1857, vol. X, plate I, fig. 6, fide Fischer-Piette, manuscript

Mediterranean record (Israel): BARASH & DANIN, 1982: 107

Material: Off Bardawil Lagoon, 1. IV. 1974, on the beach, 1 damaged valve, length 30 mm, height 19 mm (NS 19735)

Distribution: Philippines, Nicobar Is., Japan, New Guinea (Fischer-Piette, Ms. on Psammobiidae).

Trapeziidae

Trapezium oblongum (Linnaeus, 1758)

KAY 1979: 566, fig. 184 E, F

syn. *Chama oblonga* Linnaeus, 1758; DODGE 1952: 137

Mediterranean record (Israel): MIENIS 1980c: 313

Material: Rafah, IV. 1980, on the beach, 1 valve, length 43 mm, height 25 mm (NS 19736)

Distribution: Gulf of Suez (LAMY 1938: 25); Gulf of Aqaba (Dafni, pers. comm.); Red Sea, Perim (LAMY 1916b: 315); Djibouti (MASTALLER 1979: 152); Madagascar (FRANC 1960: 2099); Seychelles, Reunion Is. (LAMY 1918-19: 271); Mauritius (VIADER 1937: 64); Cocos Keeling Is. (ABBOTT 1950: 95); Japan (KIRA 1968: 148); New Guinea-Aroe Is., Tahiti, Tuamotu (LAMY 1918-19: 271); Hawaii Is. (KAY 1979: 566).

Veneridae

Gafrarium callipygum (Born, 1780)

FISCHER-PIETTE & VUKADINOVIC 1975: 35

syn. *Cytherea callipyga* (Born, 1780) ISSEL 1869: 66

Circe callipyga (Born)

Venus callipyga Born, 1780

Cytherea arabica (Lamarck, 1818); FISCHER-PIETTE & VUKADINOVIC 1975: 35, 36

Lioconcha arabica (Lamarck, 1818)

Gafrarium arabicum (Lamarck, 1818); SPRY 1964: 32

A record new to the Mediterranean (Det. Metivier)

Material: Tel Aviv, 19. X. 1980, on the beach, 1 valve, (NS 19733)

Distribution: Suez Canal, Lake Timsah, Great and Little Bitter Lakes (MOAZZO 1939: 80); Gulfs of Suez and Aqaba, Red Sea proper (STURANY 1899: 26 [280], 39 [293]); Persian Gulf, Arabian Sea-Karachi (MELVILL & STAN- DEN 1906: 830).

Order Myoida

Myidae

Sphenia rueppelli A. Adams, 1850

MOAZZO 1939: 106, fig. 5

syn. *Cuspidaria adenensis* Jousseume, 1888; LAMY 1926: 145

A record new to the Mediterranean (Israel) – (Det. Morris-Whybrow)

Material: Netanya, 14. II. 1978, on the beach, under stone, 1 shell, length 10.90 mm, height 7.35 mm, width 6.25 mm. (Registration number of British Museum – BM/NH 1980 11/8)

Distribution: Suez Canal, Port Taufiq, Little Bitter Lake (Moazzo, 1939: 106, 265); Gulf of Suez (MCANDREW 1870: 445); Red Sea, Aden, Perim, Djibouti (LAMY 1926: 145)

Remarks: Dr. Solene Morris-Whybrow, British Museum (Natural History) stated in a letter of 1. VII. 1980 "The specimen found in Israel was identified after comparison with the holotype of *Sphenia ruppelli* A. Adams, 1851, collected from the Red Sea (Proc. Zool. Soc. London, 1850: 89) Dr. Jousseume described this species in 1888: 202 under the name *Cuspidaria adenensis*. In his notes in the manuscript he remarked: "This species lives within madreporic rocks and is subjected to numerous deformations. . . . The posterior end es short and wide in most specimens, and is often rather long and narrow as in the *Cuspidaria*, therefore I considered it as a *Cuspidaria* (LAMY 1926: 146).

Gastrochaenidae

Gastrochaena cymbium Spengler, 1783*

MOAZZO 1939: 107

Mediterranean records (Israel): BARASH & DANIN 1973: 342, fig. 24; 1977: 102

(Italy): ANONYMOUS 1980: 20

Material: 1) Israel – data on material collected, see BARASH & DANIN 1973: 342; 1977: 102

2) At Talamone (central Tyrhenian Sea), 1 dead specimen has been found, III. 1979, on the beach. Capsule was attached to the interior of a valve of *Venus verrucosa*, size of capsule: height 12 mm, width 6.5 mm (Dr. Nofroni, Roma, in litt.)

General Remarks

The present report includes, as mentioned in the introduction, 29 Indo-Pacific species found in the Mediterranean, which were not discussed in the earlier reports of 1948–1977. The first records on the presence in the Mediterranean and on their distribution in this sea are indicated in Table I. Of these species 19 belong to Gastropoda (Prosobranchia – 11, Opisthobranchia – 8) and 10 to Bivalvia (Pteriomorphia – 4, Heterodonta – 7). The majority of the species (23) have been collected only along the Mediterranean coasts of North Sinai and Israel, close to the Suez Canal.

The 4 following species are recorded only from Mediterranean sites outside of Israel and Sinai: *Clelandella infuscata* – Sicily; *Nerita sanguinolenta* and *Scapharca demiri* – Aegean Sea and *Rapana venosa* – Black Sea, Adriatic Sea, Tyrrhenian Sea. 2 species: *Stombus decorus* and *Palmadusta gracilis* were found both in Israel and in the Aegean Sea (Turkey).

With regard to the vertical distribution 14 species have been collected in the intertidal zone or washed ashore, and the equal number of species (14) in the infralittoral zone, in depths of 1.5–90 m.

Shells of one species *Soletellina rubra* were obtained both in the intertidal and the infralittoral zones. Live specimens were found of the following 10 species: *Strombus decorus*, *Palmadusta gracilis*, *Rapana venosa*, *Kleinella fulva*, ?*Monoptygma sinuata*, *Plocamopherus ocellatus*, *Dendrodoris nigra*, *Acar plicatum*, *Scapharca demiri* and *Scapharca inaequivalvis*. These species are obviously able to adapt themselves to the changed ecological conditions existing in the Mediterranean.

Supplementary data are presented in this report for 4 species already discussed in the former reports: *Trochus erythraeus*, *Bursatella leachi savigniana*, *Hypselodoris infucata* (= *Glossodoris runcinata*) and *Gastrochaena cymbium*.

Only one shell of *Trochus erythraeus* was found at first (BARASH & DANIN 1973: 305) but recently 2 live specimens of this species were reported. Thus, *T. erythraeus* may probably be considered as a rock dweller in the Mediterranean waters of Israel.

Bursatella leachi savigniana was distinguished as a successful immigrant distributed in the Eastern and Western Mediterranean (BARASH & DANIN 1982: 108). The present report offers additional data on the occurrence of this species in two islands of the Aegean Sea: Khios and Lesbos.

Table 1. List of Indo-Pacific Species of Mollusca in the Mediterranean not discussed in earlier report (1948–1977)

Species	First record of presence in the Mediterranean	Distribution in the Mediterranean
Gastropoda		
Prosobranchia		
<i>Clelandella infuscata</i> (Gould, 1861)	DI NATALE 1982: 573	Sicily-Straits of Messina
<i>Nerita sanguinolenta</i> Menke, 1829	NORDSIECK 1973: 4	Karpathos Is. (Aegean Sea)
<i>Planaxis punctostriatus</i> Smith, 1872	MIENIS 1981: 361	Dor (Israel)
<i>Cerithium nesioticum</i> Pilsbry & Vanatta, 1906	MIENIS 1977 a: 45	Shiqmona (Israel)
<i>Cerithiopsis pulvis</i> (Issel, 1869)	AARTSEN & CARROZZA 1983: 37	Haifa, Hadera (Israel)
<i>Hipponix conicus</i> (Schumacher, 1817)	Present paper	Dor (Israel)
<i>Strombus decorus</i> (Roeding, 1798)	MASSILIA 1983: 21	Turkey, Rhodes, Cyprus, Israel
<i>Erosaria turdus</i> (Lamarck, 1810)	Present paper	Dor (Israel)
<i>Palmadusta gracilis</i> (Gascoïn, 1849)	BLOECHER 1983: 26	Israel: Akhziv, Nahriyya, Akko, Hadera, Palmahim, Aegean Sea – Turkey: between Mersin and Silifk Black Sea, Adriatic Sea, Tyrrhenian Sea
<i>Rapana venosa</i> (Valenciennes, 1846)	STARK 1950 (fide MIENIS 1976 a: 39)	North Sinai, near Bardawil Lagoon
<i>Vasum turbinellus</i> (Linnaeus, 1758)	MIENIS 1973: 6	
Opisthobranchia		
<i>Chrysallida fischeri</i> (Hornung & Mermod, 1925)	AARTSEN & CARROZZA 1979: 29	Haifa Bay (Israel)
<i>Kleinella fulva</i> (A. Adams, 1851)	LAVALLEYE & BARASH 1981: 87	Israel: Haifa Bay, Atlit-Dor
? <i>Monopygma sinuata</i> Gould, 1861	Present paper	Haifa bay (Israel)
<i>Cingulina isseli</i> (Tryon, 1886)	AARTSEN & CARROZZA 1983: 38	Haifa bay (Israel)
<i>Bulla ampulla</i> Linnaeus, 1758	BARASH & DANIN 1982: 107	Gaash – north of Tel Aviv (Israel)
<i>Ventommestia girardi</i> (Audouin, 1827)	MIENIS, 1976 b: 209	North Sinai, near Bardawil Lagoon
<i>Plocamopherus ocellatus</i> Rueppell & Leuckart, 1828	BARASH & DANIN 1982 a: 107	Nizanim (Israel)
<i>Dendrodoris nigra</i> Stimpson, 1855	Present paper	Israel: Dor, Caesarea
Bivalvia		
Pteriomorpha		
<i>Acar plicatum</i> (Dillwyn, 1871)	Present paper	Netanya (Israel)
<i>Scapharca demiri</i> Piani, 1981	DEMIR, 1977: 197	Aegean Sea – Turkey: Izmir
<i>Scapharca inaequalis</i> (Bruguïere, 1789)	GHSOTTI, 1973: 68	Adriatic Sea
Heterodonta		
<i>Divaricella angulifera</i> (von Martens, 1880)	MIENIS, 1979: 200	Israel: Netanya, off Bardawil (North Sinai)
<i>Laevicardium flavum</i> (Linnaeus, 1758)	Present paper	Dor (Israel)
<i>Soletellina rubra</i> (Schroeter, 1788)	MIENIS, 1980 b: 301	Israel: North Sinai – Bardawil
<i>Soletellina subradiata</i> (Reeve, 1857)	BARASH & DANIN 1982 a: 107	North Sinai – Bardawil
<i>Trapezium oblongum</i> (Linnaeus, 1758)	MIENIS, 1980 c: 313	Rafa (north of North Sinai)
<i>Gafrarium callipygum</i> (Born, 1780)	Present paper	Tel Aviv (Israel)
<i>Sphenia rueppelli</i> (A. Adams, 1850)	Present paper	Netanya (Israel)

Hypselodoris infucata was first mentioned in the report of BARASH & DANIN 1977: 98 under the name *Glossodoris runcinata*. Then, only two specimens were found. Today this species is fairly common in some localities along the Mediterranean coast of Israel. The new records are given (MIENIS & GAT 1981: 51, GAT & FAIZILBER 1983: 17) under the name *Hypselodoris infucata*, which is synonymous with *Glossodoris runcinata*. This species has apparently established itself as a permanent settler in the South Eastern Mediterranean.

Gastrochaena cymbium was reported as an immigrant into the Mediterranean, but only from the coasts of North Sinai and Israel. Dr. Nafroni (Italy) acknowledged finding shells in Talamone (Central Tyrrhenian Sea). (ANONYMOUS, 1980: 20, Table 15.) The distribution of *G. cymbium*, in the Mediterranean, outside of Israel, still needs further clarification.

In conclusion, it would be expedient to make some remarks on all of the Indo-Pacific immigrants (Lessepsian immigrants) into the Mediterranean. Altogether 91²) species of Mollusca from the Indo-West Pacific region are known to be found in the Mediterranean. They belong to 3 classes: Polyplacophora – 1 species, Gastropoda – 57, Bivalvia – 33 species. It seems that classes confined exclusively to marine environments are less capable (Polyplacophora) or incapable (Scaphopoda, Cephalopoda) of adapting themselves to the change of ecological conditions involved in the migration.

The provenance of the species that have immigrated into the Mediterranean is mostly designated in general terms: Red Sea, Indo-Pacific or Indo-West-Pacific species. An attempt should be made here to mark the subdivisions in the Indo-Pacific from which the immigrants arrived into the Mediterranean. The following subregions may be tentatively suggested: Erythraean, Western Indian Ocean (East Africa and the adjacent islands), Persian, Indian (South and East Indies), Indo-Malayan, West Pacific Is. (including Hawaii Is.) and North Australia. The subdivisions proposed are close to the schemes outlined by SCHILDER 1965; EKMAN 1967 and EALES 1960.

Table 2. Distribution of the Lessepsian immigrants in the subregions of the Indo-Pacific

Abbreviations:

SC	Suez Canal
Ery.	Erythraean subregion
WIO	Western Indian Ocean
Per.	Persian subregion
Ind.	Indian (South and East Indies) subregion
Mal.	Indo-Malayan subregion
WPI	Western Pacific Islands
Au	Australia

Species	SC	Ery.	WIO	Per.	Ind.	Mal.	WPI	Au
Polyplacophora								
<i>Chiton platei</i> Thiele, 1910		+	+		+	+		
Gastropoda								
<i>Haliotis pustulata cruenta</i> Reeve, 1846		+	+					+
<i>Diodora rueppelli</i> (Sowerby, 1834)	SC	+	+	+				+
<i>Cellana radiata</i> (Born, 1778)	SC	+	+	+	+	+		+
<i>Clelandella infuscata</i> (Gould, 1861)								+

²) 3 additional species have been recorded by Mienis (1984: 3): *Diloma atrovirens*, *Cypraea pantherina* (Lightfoot, 1786) and *Nassarius glans* (Linnaeus, 1758). The data are incomplete. These species are no doubt occasional migrants that arrived on Mediterranean beaches, probably mainly as ballast.

<i>Trochus erythraeus</i> Brocchi, 1823	SC	+	+	+			
<i>Minolia nedyma</i> Melvill, 1897		+		+	+		
<i>Umbonium vestiarium</i> (Linnaeus, 1758)		+		+		+	
<i>Nerita sanguinolenta</i> Menke, 1829	SC	+					
<i>Alvania dorbignyi</i> (Andouin, 1827)	SC	+					
<i>Rissoina bertholleti</i> (Issel, 1869)	SC	+	+	+			
<i>Planaxis punctostriatus</i> Smith, 1872		+	+			+	+
= <i>Planaxis lineolatus</i> Gould, 1849							
<i>Pirenella cailliaudi</i> (Potiez & Michaud, 1838)	SC	+	+				
<i>Bittium proteum</i> (Jousseume, 1930)		+					
= <i>Dahlakia</i> cf. <i>leilae</i> Biggs, 1971							
<i>Eufenella pupoides</i> (A. Adams, 1860)	SC	+		+	+	+	
<i>Clathrofenella reticulata</i> (A. Adams, 1860)		+		+		+	
<i>Scaliola</i> cf. <i>elata</i> Issel, 1869	SC	+	+	+			
<i>Diala varia</i> A. Adams, 1861	SC	+	+		+	+	+
<i>Cerithium erythraeonense</i> Lamarck, 1822	SC	+	+				
<i>Cerithium scabridum</i> Philippi, 1849	SC	+		+	+		
<i>Cerithium nesioticum</i> Pilsbry & Vanatta, 1906		+	+		+	+	+
<i>Rhinoclavis kochi</i> (Philippi, 1848)	SC	+	+	+		+	
<i>Cerithiopsis pulvis</i> (Issel, 1869)		+					
<i>Hipponix conicus</i> (Schumacher, 1817)		+	+			+	+
<i>Strombus lentiginosus</i> Linnaeus, 1758			+		+	+	+
<i>Strombus decorus</i> (Roeding, 1798)			+	+	+	+	
<i>Erosaria turdus</i> (Lamarck, 1810)	?SC	+	+	+	+	+	
<i>Erronea caurica</i> (Linnaeus, 1758)		+	+	+	+	+	+
<i>Palmadusta gracilis</i> (Gascoïn, 1849)		+	+	+	+	+	+
<i>Monetaria moneta</i> Linnaeus, 1758		+	+	+	+	+	+
<i>Murex tribulus</i> (Linnaeus, 1758)	SC	+	+	+	+		
<i>Aspella anceps</i> (Lamarck, 1822)		+	+	+		+	+
<i>Rapana rapiformis</i> (Born, 1778)		+	+	+	+	+	
<i>Rapana venosa</i> (Valenciennes, 1846)						+	
<i>Thais carinifera</i> (Lamarck, 1822)	SC	+	+	+	+	+	
<i>Quoyula madreporarum</i> (Sowerby, 1832)		+	+		+	+	+
<i>Anachis savignyi</i> (Moazzo, 1939)		+					
<i>Nassarius arcularius plicatus</i> (Roeding, 1798)		+	+	+			
<i>Fusinus marmoratus</i> (Philippi, 1846)	SC	+	+	+			
<i>Vasum turbinellus</i> (Linnaeus, 1758)		+	+		+	+	+
<i>Lophiotoma indica</i> (Roeding, 1798)					+	+	+
<i>Conus arenatus</i> Hwass in Bruguiere, 1792	SC	+	+		+	+	
<i>Chrysallida fischeri</i> (Hornung & Mermod, 1925)		+					
<i>Chrysallida maiae</i> (Hornung & Mermod, 1924)		+					
<i>Kleinella fulva</i> (A. Adams, 1851)				+	+	+	
? <i>Monoptygma sinuata</i> Gould, 1861						+	
<i>Cingulina isseli</i> (Tryon, 1886)		+					
<i>Ventomnestia girardi</i> (Audouin, 1827)	SC	+				+	
<i>Bulla ampulla</i> Linnaeus, 1758		+	+	+	+	+	+
<i>Notarchus indicus</i> Schweigger, 1820	?SC	+	+		+	+	+
<i>Bursatella leachi savigniana</i> Audouin, 1826	SC	+					
<i>Pleurobranchus forskali</i> Rueppell & Leuckart, 1828		+	+				
<i>Berthellina citrina</i> (Rueppell & Leuckart, 1828)	SC	+	+		+	+	+
<i>Plocamopherus ocellatus</i> Rueppell & Leuckart, 1828	SC	+					
<i>Hypselodoris infucata</i> (Rueppell & Leuckart, 1828)		+				+	+
= <i>Glossodoris runcinata</i> (Bergh, 1877)							
<i>Discodoris concinna</i> (Alder & Hancock, 1864)		+	+		+		+
<i>Dendrodoris nigra</i> Stimpson, 1855	SC	+	+	+	+	+	+
<i>Siphonaria kurracheensis</i> Reeve, 1856	SC	+		+			+

Species	SC	Ery.	WIO	Per.	Ind.	Mal.	WPI	Au
Bivalvia								
<i>Acar plicatum</i> (Dillwyn, 1817)	SC	+	+	+		+	+	+
<i>Scapharca demiri</i> Piani, 1981 = <i>Arca amygdalum</i> Philippi, 1847						+		
<i>Scapharca inaequivalvis</i> (Bruguere, 1789)						+		
<i>Scapharca natalensis</i> (Krauss, 1848)	SC	+	+	+		+		
<i>Limopsis multistriata</i> (Forskall, 1775)		+	+		+			
<i>Glycymeris</i> cf. <i>arabica</i> (H. Adams, 1870)	SC	+						
<i>Modiolus auriculatus</i> Krauss, 1848	SC	+	+	+		+	+	+
<i>Modiolus arcuatulus</i> Hanley, 1843		+	+	+		+	+	
<i>Modiolus glaberrimus</i> (Dunker, 1856)	SC	+	+			+		+
<i>Brachidontes variabilis</i> (Krauss, 1848)	SC	+	+		+		+	+
<i>Pinctada radiata</i> (Leach, 1814)	SC	+	+	+	+	+	+	+
<i>Malleus regula</i> (Forskall, 1775)	SC	+	+	+		+	+	+
<i>Spondylus spectrum</i> Reeve, 1856		+	+			+		
<i>Crassostrea</i> cf. <i>gigas</i> (Thunberg, 1793)						+	+	
<i>Divaricella angulifera</i> (von Martens, 1880)	SC	+	+			+		
<i>Chama broderipi</i> Reeve, 1846	SC		+			+		
<i>Pseudochama cornucopia</i> (Reeve, 1846)	SC	+						
<i>Laevicardium flavum</i> (Linnaeus, 1758)		+	+	+	+	+	+	+
<i>Papyridea australe</i> (Sowerby, 1834)	SC	+	+	+		+	+	+
<i>Papyridea papyraceum</i> (Gmelin, 1791)	SC	+	+	+	+	+	+	+
<i>Hippopus hippopus</i> (Linnaeus, 1758)						+	+	+
<i>Macraolorina</i> Philippi, 1846	SC	+	+	+		+		
<i>Atactodea striata</i> (Gmelin, 1791)	SC	+	+			+	+	
<i>Angulus valtonis</i> (Hanley, 1844)	SC	+						
<i>Soletellina rubra</i> (Schroeter, 1788)	SC	+	+	+				
<i>Soletellina subradiata</i> (Reeve, 1857)						+		
<i>Trapezium oblongum</i> (Linnaeus, 1758)		+	+		+	+	+	
<i>Gafrarium callipygum</i> (Born, 1780)	SC	+	+	+				
<i>Garfarium pectinatum</i> (Linnaeus, 1758)	SC	+	+	+	+	+	+	
<i>Clementia papyracea</i> (Gmelin, 1791)	SC	+		+	+	+		+
<i>Paphia textile</i> (Gmelin, 1791)	SC	+		+	+	+		+
<i>Sphenia rueppelli</i> A. Adams, 1850	SC	+						
<i>Gastrochaena cymbium</i> Spengler, 1783	SC	+			+	+		
<i>Laternula subrostrata</i> (Lamarck, 1818)	SC	+	+	+	+	+	+	

Table 3. Distribution of the Lessepsian immigrants in the subregions of the Indo-Pacific³⁾ (91 species = 100 %)

Erythraean subregion	79 species	86.8 %
Western Indian Ocean	56 species	61.5 %
Persian subregion	41 species	45 %
Central Indian Ocean	37 species	40 %
Indo Malayan subregion	60 species	66 %
Pacific Is.	36 species	39 %
Australia	23 species	25.3 %

Tables 2, 3, show that the greatest part of the Indo-Pacific species in the Mediterranean – 79 species – are from the Erythraean subregion: – 14 species, from the Erythraean subregion only, 65 species also from other subregions.

³⁾ *Crassostrea* cf. *gigas* (Thunberg, 1793) was recorded as an Indo-Pacific immigrant into the Mediterranean (GHISOTTI 1974: 14). It proved to be an introduction of an edible clam and not taken into account in the percentages of the distribution in the subregions of the Indo-Pacific.

A relatively small part of migrants – 12 species – are not recorded from the Erythraean subregion, but from the areas more remote from the Mediterranean Sea (Tables 3, 4).

Table 4. Lessepsian migrants not recorded from the Erythraean subregion

Species	Collected	Distribution outside the Mediterranean ⁴⁾
<i>Clelandella infusata</i> (Gould, 1861)	One shell	Japanese Sea
<i>Strombus lentiginosus</i> Linnaeus, 1758	? shells	East Africa, Madagascar, Ceylon, Indonesia, Western Pacific Is., Philippines
<i>Strombus decorus</i> (Roeding, 1798)	Live	Zanzibar, Mozambique, South Africa, Madagascar, Gulf of Aden, Persian Gulf, Saudi Arabia, Ceylon, Burma, Thailand, Indonesia
<i>Rapana venosa</i> (Valenciennes, 1846)	Live	Japan – Tartari Canal, Yellow Sea
<i>Lophiotoma indica</i> (Roeding, 1798)	? Live	Ceylon, Thailand, Malacca, Indonesia, (Java, Celebes), Philippines, Japan, New Guinea, Australia
<i>Kleinella fulva</i> (A. Adams, 1851)	Live	Arabian Sea: Karachi, Bombay, Philippines, Japan
? <i>Monoptygma sinuata</i> Gould, 1861	Live	China Seas
<i>Scapharca demiri</i> Piani, 1981	Live	China Seas
<i>Scapharca inaequivalvis</i> Bruguière, 1789	Live	Philippines, Japan
<i>Chama broderipi</i> (Reeve, 1846)	shells	Madagascar, New Guinea
<i>Hippopus hippopus</i> (Linnaeus, 1758)	One valve	Malaya, Indonesia, Philippines, Japan, Western Pacific Is., Australia
<i>Soletellina subradiata</i> (Reeve, 1857)	One valve	Nicobar Is., Philippines, Japan

It may be surmised that the pelagic larvae of the benthic molluscan species living in the Erythraean subregion are brought into the Mediterranean by sea currents and settle there. It is very unlikely that the species from subregions that are remote from the Mediterranean should be able to migrate by means of the larval stage, since the life span of the larvae is rather limited (up to 9 weeks in Gastropoda and 5 weeks in Bivalvia). It is more reasonable to assume that the migrants from the distant subregions were transported in the adult stage by carriers (ships, fish, etc.) to the localities in which they were collected in the Mediterranean.

Many Indo-Pacific species, 43 of the 91, occurring in the Mediterranean have not been recorded from the Suez Canal (see Table 2). Presumably they did not find suitable environmental conditions for settling in the Suez Canal.

The inclusion of the following species in the list of the Lessepsian migrants into the Mediterranean is questionable: *Scaliola* cf. *elata*, *Strombus lentiginosus*, *Monetaria moneta*, *Aspella anceps*, *Spondylus spectrum*, *Crassostrea* cf. *gigas* and *Hippopus hippopus*.

Scaliola cf. *elata* is recorded by BARASH & DANIN, 1977: 91, fig. 6. According to Mienis (in litt.) there are “differences between true *Scaliola* and the figured shell. It is evident that the latter is not congeneric”.

Shells of *Strombus lentiginosus* and *Spondylus spectrum* were collected on the Israeli Mediterranean coast by AHARONI (1934). No information was given about the number of specimens collected, nor about the site where they were deposited.

The appearance of *Monetaria moneta* and *Hippopus hippopus* in the Mediterranean seems to be adventitious. Their shells were most probably brought to the Israeli coast by man.

The identity of *Aspella anceps* is disputed by Vokes (in litt.): “I doubt that the species, you have, is really *anceps*, more likely it is *producta*, which does occur along the coast of eastern Africa and Mauritius. Another possibility is that the species is a native Mediterranean form that has just been overloo-

⁴⁾ The geographical records according to BARASH & DANIN, 1973 and the present report.

ked.” RADWIN & D’ATTILIO (1976: 21) also contest the identity of *Aspella anceps*, saying: “The name is probably best considered a ‘nomen dubium’.”

Crassostrea cf. *gigas* (Thunberg, 1793) was recorded as a species of Indo-Pacific origin, found in the Mediterranean (GHISOTTI 1971: 13; 1974b: 14). It seems, however, to be a case of introduction of edible clams for commercial reasons. For the past 5 years *C. gigas* has frequently been reported as being from Mediterranean and Black Sea localities.

The results of the observations made for about 40 years on the migration of Indo-Pacific Mollusca into the Mediterranean will be summarised below. The number of Indo-Pacific species reported to occur in the Mediterranean have increased fivefold (from 20 to 91) from 1948 to 1984 (Table 5).

Table 5.

	Polyplacophora	Gastropoda	Bivalvia	Total
Report HAAS, 1948	—	8	10	18
Present Report, 1984	1	57	33	91

Of the 91 species appearing in the Mediterranean, 44 species have been collected as one or a few shells, their being inhabitants of this sea is doubtful. The other 47 species were found live or in considerable numbers of shells. These species are considered as residents in the Mediterranean.

The following 13 species may safely be qualified as permanent inhabitants of the Mediterranean: *Diodora rueppelli*, *Minolia nedyma*, *Pirenella cailliaudi*⁵⁾, *Cerithium scabridum*, *Rhinoclavis kochi*, *Thais carinifera*, *Bursatella leachi savigniana*, *Hypselodoris infucata*, *Brachidontes variabilis*, *Pinctada radiata*, *Malleus regula*, *Paphia textile* and *Gastrochaena cymbium*.

We have had the possibility of following the rate of colonization of 3 successful immigrants from the beginning of their penetration into the new area. *Minolia nedyma* was recorded first by BARASH & DANIN (1973: 305) under the name *Isanda* cf. *holdsworthiana* (NEVILL, 1871). The first two specimens were dredged in 1966 at Haifa Bay. This species is now common in various localities on the continental shelf of Israel. *Rhinoclavis kochi* was recorded for the first time in the Mediterranean in 1963. It is at present one of the most abundant species in the continental shelf of Israel. *Hypselodoris infucata* was first recorded from Israel in 1974 (BARASH & DANIN, 1977). Only 2 specimens were found. In the present report the number of specimens already observed amounts to 48.

In retrospect, on the data accumulated since the opening of the Suez Canal (1869) the following may be stated on the phenomenon of the Lessepsian migration of Mollusca:

1. The number of immigrating species has increased gradually during the past years.
2. The great majority of the Indo-Pacific immigrants are concentrated close to the Suez Canal. Their penetration into other parts of the Mediterranean is increasing, but at a very slow rate.

Addendum

Further findings of *Strombus decorus* in additional localities reported by K. NICOLAY (La Conchiglia, 1986 No. 202, 203: 20): Cyprus-Aghia Napa, leg T. Zambakidis and Rhodes in Ixia by Dr. R. de Roover from Antwerpen.

Countless specimens of *Melibe fimbriata* Alder and Hancock, 1864 (Tethyidae) were observed in the Astacos inlet of the Ionian Sea coast of Greece on 22–24 September, 1982 (THOMPSON and CRAMPTON 1984: 114). This species is of Indo-Pacific origin and the number of Lessepsian immigrants of Mollusca in the Mediterranean amounts with it to 92.

⁵⁾ *Pirenella cailliaudi* considered as an Indo-Pacific species and *P. conica* – as a Mediterranean one, seem to belong to the same species.

Acknowledgements

The authors are grateful to Mr. H. K. Mienis for the information concerning the Indo-Pacific immigrants dealt with in the paper. We are much indebted to the late Dr. I. Yaron for the reading of the manuscript and his constructive remarks. His tragic death was a great loss for the Malacology of Israel.

Literature

- AARTSEN VAN, J. J. and F. CARROZZA 1979: *Chrysallida fischeri* (Hornung and Mermod, 1925): A Red Sea species found at the Israeli Mediterranean coast. – Boll. Malacologico (Conchiglie), XV (1–2): 29–30
- — 1983: Two more Red Sea species recorded for the first time from the Mediterranean coast of Israel. – Boll. Malacologico, Milano, (1–4): 37–40
- ABBOTT, R. T. 1950: The molluscan fauna of the Cocos-Keeling Islands, Indian Ocean. – Bull. of the Raffles Museum, Singapore, 22: 68–98
- — 1958: Marine Mollusca of Rennell Islands, Solomon Islands. Nat.-Hist. Rennell Isl. Br. Solomon Isl., 2: 203–206
- — 1960: The genus *Strombus* in the Indo-Pacific. – Indo-Pacific Mollusca, 1(2): 33–174
- — 1974: American seashells. – Van Nostrand Reinhold Co., New York. 663 pp., 6415 text figs., 24 coloured plates.
- ADAM, W. & E. LELOUP 1938: Prosobranchia et Opisthobranchia. – Result. Sci. Voyage Indes Orient. Nederl., II, fasc. 19, 209 pp., 8 plates.
- — 1939: Gastropoda, Pulmonata, Scaphopoda et Bivalvia – Result. Sci. Voyages Indes Orient. Nederl., II, fasc. 20, 126 pp., 7 plates.
- AHARONI, J. 1934: From Ashkelon to Rubin – Nature and Country, 2: 473–476 (in Hebrew).
- ALLAN, J. 1950: Australian shells – Georgian House, Melbourne, 470 pp. with text figures.
- ANONYMOUS 1980: Synoptic tables of European and Mediterranean shells. – La Conchiglia (The shell), XII (136–137): 20, table 15.
- BARASH, AL. & Z. DANIN 1973: The Indo-Pacific species of mollusca in the Mediterranean and notes on a collection from the Suez Canal. – Isr. J. Zool., 21(3–4): 301–374, 25 figs.
- — 1977: Additions to the knowledge of Indo-Pacific Molluscs in the Mediterranean. – Conchiglie, Milano, 13: 85–116, 15 figs.
- — 1982: Mediterranean Mollusca of Israel and Sinai: Composition and distribution. – Isr. J. Zool., 33: 86–118
- BARNARD, K. H. 1964: Contribution to the knowledge of South African marine Mollusca, part V: Lamellibranchiata. – Ann. South Afr. Mus. Cape Town, XLVII: 361–593
- BERTSCH, H. & S. JOHNSON 1979: Three new opisthobranch records for the Hawaiian Islands. – The Veliger, 22(1): 41–44, 2 figs.
- BERTSCH, H. & S. JOHNSON 1981: Hawaiian nudibranchs – Oriental Publishing Co., Honolulu, Hawaii, 112 pp. with coloured figs.
- BIGGS, H. E. J. 1969: Marine Mollusca of Masirah Is. – South Arabia. – Arch. Moll. 99(3–4): 201–207
- — 1973: The marine mollusca of the Trucial Coast, Persian Gulf. – Bull. Brit. Mus. (Nat. Hist.), London, 24(8): 343–421, 6 plates.
- BLOECHER, M. 1983: *Cypraea gracilis notata* Gill, 1858 en Mediterranee? – Xenophora, 14: 26, Paris.
- — 1983: In: (Nicolay, K. and Manoja, E. R. eds.) Recent finds. – La Conchiglia, XV (166–167): 17–18
- — 1984: New species in the Mediterranean: 2 reports. – Hawaiian Shell News, 32(2): 3
- BOUCHET, PH. & F. DANRIGAL 1982: Napoleon's Egyptian campaign (1798–1801) and the Savigny collection of shells. – The Nautilus, 96(1): –23, 101 figs.
- BURGESS, C. M. 1970: The living Cowries. – A. S. Barnes and Co., New York, 389 pp., 44 plates.
- CATALANO, E., N. PARINELLO & G. DELEO 1978: Nuovi dati sulla penetrazione nel Mar Mediterraneo di *Bursatella leachii savigniana* Audouin, 1826. Ritrovamento nel Golfo di Palermo. – Mem. Biol. Marina e Ocean., 8(5): 123–128
- CATE, C. N. 1968: Western Australian Cowries. A second revisal and expanded report, plates 21–34, 5 maps. – The Veliger, 10(3): 212–232
- CERNOHORSKY, W. O. 1967: Marine shells of the Pacific. – Pacific Publications, Sydney, 248 pp., 444 figs.
- — 1968: Observations on *Hipponix conicus* (Schumacher, 1817) The Veliger, 10(3): 275–280

- — Marine shells of the Pacific, vol. II – Pacific Publications, Sydney. 411 pp., 68 plates.
- CHENU, LE J. C. 1859; 1862: Manuel de Conchyliologie et de Paleontologie conchyliologique. – Paris Librairie Victor Masson: Vol. 1 – 1859, Gasteropodes, 508 pp., 3707 figs; Vol. 2 – 1862, Acephales, 327 pp., 1236 figs.
- COWAN, J. MC. 1974: The West American Hipponicidae and the application of *Malluivium*, *Antisabia* and *Hippornix* as generic names. – The Veliger, **16**(4): 377–380, 24 figs.
- DAUTZENBERG, PH. 1929: Mollusques testaces marins de Madagascar – Faune Colon. Fr. III: 115–430 (321–636), plates IV–VII.
- DAUTZENBERG, PH. & J. L. BOUGE 1933: Les mollusques testaces marins des etablissements Francais de l'Oceanie – J. Conchyliol., **LXXVII**: 43–371
- DEMIR, M. 1977: On the presence of *Arca* (*Scapharca*) *amygdalum* Philippi, 1847 (Mollusca, Bivalvia) in the harbour of Ismir, Turkey: Univ. Fen. Fak. Mec. Istanbul Serie B (1977), **42**: 197–202, 1 plate.
- DI GERONIMO, J. & E. ROBBA 1979: Contributo alla conoscenza della malacofauna del Benadir (Somalia Meridionale). – Boll. Malacologico, Milano, **15**(7–8): 165–196
- DI NATALE, A. 1982: Extra Mediterranean species of Mollusca along the southern Italian coasts. – Malacologia, **22**(1–2): 571–580
- DODGE, H. 1952: A historical review of the molluscs of Linnaeus, part I: The classes Loricata and Pelecypoda. – Bull. Amer. Mus. Nat. Hist. New York, vol. **100**, 263 pp.
- EALLES, N. B. 1960: Revision of the world species of *Aplysia* (Gastropoda, Opisthobranchia). – Bull. Brit. Mus. Nat. Hist. (Zool.), **5**: 269–404
- EKMAN, S. 1967: Zoogeography of the sea – Sidgewick and Jackson Ltd., London, 417 pp.
- ENGEL H. and C. J. VAN EEKEN, 1962: Red Sea Opisthobranchia from the coast of Israel and Sinai: Sea Fish. Res. Stn., Haifa Bull no **30**: 15–34
- FISCHER P. H., 1970, Gastropodes Testaces marins du Golfe de Siam. J. Conchyliol. CVIII (4): 93–121
- FRANC, A. 1960: Classe des bivalves – Traite de Zoologie (ed. Grasse) Paris, V, II fasc: 1845–2133
- GAT, G. & M. FAINZILBER 1983: Immigrants from the Red Sea – La Conchiglia, 172–173: 17
- GHISOTTI, F. 1971a: *Rapana thomasiana* Crosse, 1861 (Gastropoda, Muricidae) nel Mar Nero. – Conchiglie, **VII**(3–4): 55–58
- — 1971b: Molluschi del genere *Crassostrea* nell'Alto Adriatico – Conchiglie, **VII**(7–8): 113–124, with figs.
- — 1972: Rinvenimenti malacologici nel Mediterraneo – Conchiglie, **VIII**(1–2): 20–21
- — 1973: *Scapharca* cfr *cornea* (Reeve) ospite nuova del Mediterraneo. – Notiziario U. M. J: 68
- — 1974a: *Rapana venosa* (Valenciennes) nuova ospite Adriatica? – Conchiglie, X(5–6): 125–126
- — 1974b: Recente penetrazione in Mediterraneo di molluschi marini di provenienza Indo-Pacifica. – Quaderni della Civica Stazione Idrobiologica di Milano, **5**: 7–16, 3 plates.
- GHISOTTI, F. & E. RINALDI 1976: Osservazione sulla popolazione di *Scapharca* insediatai in questi ultimi anni su un tratto del litorale romagnolo. – Conchiglie, **12**(9–10): 183–195
- GROSSU, A. V. & D. LUPU 1964: The presence of *Rapana bezoar* opposite the Rumanian Black Sea shores (Muricidae). – Arch. Moll., **93**(5–6): 215–218
- HAAS, G. 1948: Sur l'immigration de mollusques Indo-Pacifique dans les eaux cotiers de la Palestine. – J. Conchyliol., **LXXXVIII**: 141–144
- HABE, T. 1970 (1964): Shells of the Western Pacific, in colour. vol. II Osaka, Japan, 233 pp., 66 colour plates.
- HIDALGO, J. G. 1905: Noticia sobre las faunas malacologicas del Archipelago del Jolo e Islas Marianas. – Molluscos marinos. Rev. R. Acad. Cienc. Exact., Madrid, II(4): 3–16
- HORNUNG, A. & G. MERMOD 1925: Mollusques de la Mer Rouge recueillis par A. Issel, 2nd partie. – Ann. del Mus. Civ. St. Nat. de Genova, **52**: 20–33
- ISSEL, A. 1869: Malacologia del Mar Rosso – Pisa, 387 pp., 5 plates.
- JICKELI, C. F. 1874–1884: Studien über die Conchylien des Rothen Meeres. – Jb. Dt. Malak. Ges. I: 15–263
- JOHNSON, R. I. 1964: The recent Mollusca of Augustus Addison Gould – Smith. Inst. U., Nat. Mus., 182 pp., 45 plates.
- JOUSSEAUME, F. 1888: Description des mollusques recueillis par M. le Dr. Faurot dans la Mer Rouge et le Golfe d'Aden. – Memoires de la Societe Zoologique de France: 165–223
- — 1930: Cerithiidae de la Mer Rouge – J. Conchyliol., **74**: 270–296
- KAY, A. E. 1979: Hawaiian shells – Bishop Museum Press, Honolulu, Hawaii, 633 pp., 195 figs.
- KILBURN, R. & E. RIPPEY 1982: Sea shells of southern Africa – Macmillan South Africa, Johannesburg, 249 pp., 45 coloured plates.

- KIRA, T. 1968 (1962): Shells of the Western Pacific (in colour). – Vol. I, 224 pp., 72 plates.
- KURODA, T., HABE, T. & K. OYAMA 1971: The sea shells of Sagami Bay. 489 pp., 121 plates.
- LAMY, E. 1906: Liste des Lamellibranches recueillis par M. L. G. Seurat au îls Tuamotu et Gambier (1902–1905). – Bull. Mus. Hist. Nat., 12: 205–215
- — 1907: Revision des *Arca* vivants du Museum d'Histoire Naturelle a Paris. – Journ. Conchyliol., 1–307, 3 plates.
- — 1916 a: Les lucines et les diplodontes de la Mer Rouge (d'après les matériaux recueillis par M. le Dr. Jousse-aume). – Bull. Mus. Hist. Nat., 3, 4: 1–19
- — 1916 b: Les cardites et les cypricardes de la Mer Rouge (d'après les matériaux recueillis par M. le Dr. Jousse-aume). – Bull. Mus. Hist. Nat., XXII: 311–316
- — 1917: Les arches de la Mer Rouge – Bull. Mus. Hist. Nat., 1, 2: 308–323
- — 1918: Les psammobies de la Mer Rouge (d'après les matériaux recueillis par M. le Dr. Jousse-aume). – Bull. Mus. Hist. Nat. Paris, 4: 242–250
- — 1918–1919: Revision des Cypricardiacea et des Isocardiacea vivants du Museum d'Histoire Naturelle de Paris. – J. Conchyliol.: 259–307
- — 1920: Revision des Lucinacea vivants du Museum d'Histoire Naturelle de Paris. – J. Conchyliol.: 233–318
- — 1926: Les myes de la Mer Rouge (d'après les matériaux recueillis par le Dr. Jousse-aume): 140–146
- — 1938: Mollusca testacea. Mission Robert Ph. Dollfus en Egypte. – Mem. Inst. Egypte, 37, 89 pp., 10 text figs., 1 plate.
- LAVALEYE, M. S. S. & AL. BARASH 1981: First record of the Indo-Pacific species *Kleinella (Actaeopyramis) fulva* (A. Adams, 1851) (Gastropoda, Pyrammidellidae) from the Mediterranean. – Basteria, 45(4–5): 87–89
- MARCUS, E. & E. MARCUS 1970: Some Gastropods from Madagascar and West Mexico. – Malacologia, 10(1): 181–223
- MASSILIA, L. R. 1983: Turkey: Aegean and East Mediterranean coasts. – La Conchiglia, XV (174–175): 3, 20–21
- MASTALLER, M. 1978: The marine molluscan assemblages of Port Sudan, Red Sea. – Zoologische Mededelingen, Rijksmuseum van Natuurlijke Historie Leiden, 53(13): 117–144
- — 1979: Beiträge zur Faunistik und Oekologie der Mollusken und Echinodermen in den Korallenriffen bei Aqaba, Rotes Meer – Thesis, 344 pp. Ruhr Universität, Bochum.
- MCANDREW, R. 1870: XLIII Report on the testaceous Mollusca obtained during a dredging excursion in the Gulf of Suez in the months of February and March 1869. – The Annals and Magazine of Natural History, 36: 429–450
- MEL, P. 1976: Sulla presenza di *Rapana venosa* (Valenciennes) et di *Charonia variegata* Seguenzae (Ar. e Ben.) nell'Alto Adriatico. – Conchiglie, Milano 12(5–6): 129–132
- MELVILL, J. C. & R. STANDEN 1901: The Mollusca of the Persian Gulf, Gulf of Oman and Arabian Sea, vol. II, part I, Cephalopoda, Gastropoda, Scaphopoda. – Proc. Zool. Soc. London: 327–460, 3 plates.
- — 1906: The mollusca of the Persian Gulf, Gulf of Oman and Arabian Sea, vol. VII, part II, Pelecypoda. – Proc. Zool. Soc. London: 783–848, 4 plates.
- MIENIS, H. K. 1971: Cypraeidae from the Sinai area of the Red Sea. – Argamon, 1970 1(1): 13–48
- — 1973: *Vasum turbinellus* from the Mediterranean. – Argamon, 4(1): 6
- — 1974: Notes on recent and fossil Neritidae *Neritina kinzelbachi* Nordsieck (a synonym of *N. sanguinolenta* Menke). – Argamon, 4(2–4): 45–46
- — 1976 a: On the identity of *Rapana pontica* Nordsieck, 1969 – Soosiana, 4: 39–43
- — 1976 b: *Ventomnestia girardi* (Audouin, 1827) from the Mediterranean. – Conchiglie, Milano, 12(9–10): 209–210
- — 1977 a: *Cerithium nesioticum* Pilsbry and Vanatta, 1906. Another Indo-Pacific species from the Mediterranean coast of Israel. – Conchiglie, Milano, 13(1–2): 45
- — 1977 b: A key to the Nerita species from the Sinai area of the Red Sea with various other notes. – Levantina, 7: 71–73
- — 1979: *Divaricella angulifera* (von Martens 1880). Another Red Sea species from the Mediterranean coast of Israel. – Levantina, 18: 200
- — 1980 a: *Planaxis lineolatus* Gould. Another Indo-Pacific species from the Mediterranean coast of Israel. – Levantina, 28: 324
- — 1980 b: *Soletellina rueppelliana*. Another Erythraean species from the Mediterranean coast of Israel. – Levantina, 26: 31

- — 1980 c: A record of the Indo-Pacific species *Trapezium oblongum* from the Mediterranean coast of Israel. – *Levantina*, **27**: 313–314
- — 1981: Notes concerning the nomenclature of Red Sea molluscs. 5. *Planaxis (Angiola) punctostriatus* Smith, 1872 – *Levantina*, **30**: 361–362
- — 1984: New species in the Mediterranean. Two reports. – *Hawaiian Shell News*, **32**(2): 3
- MIENIS, H. K. & M. GAT 1981: On the presence of the Indo-Pacific nudibranch *Hypselodoris infucata* along the Mediterranean coast of Israel. – *Argamon*, **7**(4): 51–52
- MIENIS, H. K. & B. SINGER 1983: An interesting addition to the mollusc fauna of the Mediterranean coast of Israel. – *Levantina*, **46**: 535–537
- MOAZZO, P. G. 1939: Mollusques testaces marins du Canal de Suez. – *Mem. Inst. Egypte, Le Caire*, **38**, 284 pp., 26 text figs, 14 plates, 1 colour plate.
- NICOLAY, K. & E. R. MANOJA 1983: *Strombus (Conomurex) decorus raybaudi* n. sp. – *La Conchiglia*, **XV**(176–177): 17–18
- NORDSIECK, F. 1973: New finds in the Mediterranean. *Neritina kinzelbachi*. – *La Conchiglia*, 7–8 (53–54): 4
- O'DONOGHUE, C. H. 1929: Report on the Opisthobranchiata, Cambridge Expedition to the Suez Canal, 1924. – *Trans. Zool. Soc. London*, **22**: 713–841
- ORE, J. 1980: Annotated checklist of Hong Kong nudibranchs. – *Hong Kong University Press*: 109–117
- ORR MAES, V. 1967: The littoral marine molluscs of Cocos-Keeling Islands (Indian Ocean). – *Proc. of the Acad. Nat. Sci. of Philadelphia*, **119**(4): 93–217, 26 plates.
- PALLARY, P. 1912: Catalogue des mollusques du littoral Mediteraneen de l'Egypte. – *Mem. Inst. Egypte*, **VII**: 200 pp., plates XV–XVIII.
- — 1926: Explication de planches de J. C. Savigny. – *Mem. Inst. Egypte, Le Caire* **XI**, 138 pp., 18 plates.
- PIANI, P. 1980: Ritrovamento nelle acque della Sicilia Orientale di *Bursatella leachii savigniana* (Audouin 1826) Opisthobranchia: Aplysiomorpha. – *Boll. Malac. Milano*, **XVI** (1–2): 5–8
- — 1981: *Scapharca demiri* nomen novum pro *Arca amygdalum* Philippi. – *Boll. Malac. Milano*, **17**(11–12): 284
- POR, F. D. 1978: Lessepsian migration – *Springer Verlag*, 228 pp., 10 plates, 48 text figs.
- PRASHAD, B. 1932: The Lamellibranchia of the Siboga Expedition. – *Leiden Bull.*, 353 pp., 9 plates.
- PRUVOT-FOL, A. 1933: Opisthobranchiata. Mission Robert Ph. Dollfus en Egypte. – *Mem. Inst. d'Egypte*: 89–160, 4 plates.
- RADWIN, G. E. & A. D'ATTILIO 1976: Murex shells of the world. An illustrated guide to the Muricidae. – *Stanford University Press*, 284 pp.
- REGTERN, ALTEA C. O. VAN 1945: Report upon a collection of recent shells from Java. – *Zool. Meded.*, **XXV**: 140–154, 2 figs.
- RINALDI, E. 1977: Primi stadi di sviluppo di *Scapharca inaequivalvis* (Bruguiera). – *Conchiglie, Milano*, **13**(11–12): 199–202
- RIPPINGALE, O. H. & D. F. MCMICHAEL 1961: Queensland and Great Barrier Reef shells: Brisbane. – *The Jacaranda Press*, 210pp., 23 coloured plates.
- ROBERTSON, R. 1969: On some molluscs collected from southwest Ceylon during the International Indian Ocean Expedition 1964. – *Extract from the Spolia Zeylanica*, **31**(II): 1–8
- RUDMAN, W. B. 1977: Chromodorid opisthobranch Mollusca from East Africa and the tropical West Pacific. – *Zoological Journal of the Linnean Society*, **61**: 351–397, 1 plate, 20 figs.
- SAFRIEL, U. & Y. LIPKIN 1964: Note on the intertidal zonation of the rocky shores at Eilat (Red Sea, Israel). – *Isr. J. Zool.*, **13**: 187–190
- SATYAMURTI, S. T. 1952: The Mollusca of Krusdai Island (in the Gulf of Manaar) I. Amphineura and Gastropoda. – *Bull. Madras Govern. Mus.*, **I**(2): 265 pp., 34 plates.
- — 1956: The Mollusca of Krusdai Island (in the Gulf of Manaar). II. Scaphopoda, Pelecypoda and Cephalopoda. – *Bull. Madras Govern. Mus.*, **I**(2): 202 pp., 30 plates.
- SCHILDER, F. A. 1956: *Lehrbuch der Allgemeinen Zoogeographie* – *Gustav Fischer Verlag, Jena*. 150 pp.
- SHOPLAND, E. R. 1902: List of marine shells collected in the neighbourhood of Aden between 1892 and 1901. – *Proc. Malacol. Soc. Lond.*, **V**: 171–179
- SMITH, E. A. 1891: On a collection of marine shells from Aden with some remarks upon the relationship of the molluscan fauna of the Red Sea and the Mediterranean. – *Proc. Zool. Soc. Lond.*, **XXVIII**: 390–445
- SMYTHE, K. R. 1972: Marine Mollusca from Bahrain Island, Persian Gulf. – *J. Conch.*, **27**: 491–496
- — 1979: The marine Mollusca of the United Arab Emirates, Arabian Gulf. – *J. Conch.*, **30**: 57–80, 11 figs.

- SOLEM, A. 1953: Marine and freshwater molluscs of the Solomon Islands. – *Fieldiana Zool.*, **34**: 213–227
- SPRY, J. F. 1961: The sea shells of Dar es Salaam. – *Tanzania Soc.*, part I, *Gastropoda*, 40 pp., 8 plates.
- — 1964: The sea shells of Dar es Salaam. Part II: *Pelecypoda (Bivalves)*, 41 pp., 8 plates.
- STARMUEHLNER, F. 1974: Beiträge zur Kenntnis der Mollusken Fauna im Littoral von Südindien und Ceylon. – *J. mar. biol. Ass. Indian*, **16**(1): 49–82
- STURANY, R. 1899: Lamellibranchiaten des Rothen Meeres. Expeditionen S. M. Schiff „Pola“ in das Rothe Meer 1895/96 und 1897/98. – *Zool. Ergeb.*, Wien, **XIV**: 1–41 (255–295), 7 plates.
- — 1903: Gastropoden des Rothen Meeres. Expeditionen S. M. Schiff „Pola“ in das Rothe Meer 1895/96, 1897/98. – *Zool. Ergeb.* Wien, **XXIII**: 1–75 (208–283)
- TAPPARONE-CANEFERI, C. 1874: Contribuzioni per una fauna malacologica delle isole Papuane. – *Ann. Mus. Civ. St. Genova*, **VI**: 21 pp.
- TAYLOR, J. & J. G. WALLS 1975: *Cowries* – T. F. H. Publications Inc. Ltd., 288 pp., with coloured figs.
- TERRENI, G. 1980: Molluschi poco conosciuti dell'Archipelago Toscano: 1. *Gastropodi*. – *Boll. Malac. Milano*, **16**, 1–2: 9–17, with figs.
- THIELE, J. 1931: *Handbuch der Systematischen Weichtierkunde*. – Gustav Fischer, Jena. 778 pp., 783 figs.
- THOMPSON, T. E. 1975: *Dorid nudibranchs from Eastern Australia (Gastropoda, Opisthoranchia)*. – *J. Zool. Lond.*, **176**: 477–517
- THOMPSON, T. E. & D. M. CRAMPTON, 1984: *Biology of Melibe fimbriata*. A conspicuous Opisthobranch mollusc of the Indian Ocean, which has now invaded the Mediterranean Sea. – *J. Moll. Stud.* **50**: 113–121
- THORSON, G. 1940: Studies on the egg masses and larval development of gastropods from the Iranian Gulf. – *Dan. Sci. Invest. Iran*, **II**: 159–238
- TINKER, S. W. 1967: *Pacific sea shells*. – Charles E. Tuttle Co, Japan, 240 pp., with figs.
- TORTORICI, R. & P. PANETTA 1977: Notizie ecologiche su aleuni Opisthobranchi raccolti nel Golfo di Taranto (*Gastropoda*). – *Atti. Soc. Ital. Sci. Nat. Museo Civ. Stor. Nat.*, Milano, **118**(2): 249–257
- TRYON, G. W. 1887: *Manual of Conchology* – Philadelphia, **IX**: 122–149, plates 19–29
- WENZ, W. 1960–62 (1938–1944): *Gastropoda*. Allgemeiner Teil und Posobranchia. In: *Handbuch der Palaeozoologie*. Berlin, **6**, 1: 1639 pp., 4211 figs.
- VIADER, R. 1937: Revised catalogue of testaceous molluscs of Mauritius and its Dependencies. – *Bull. Maurit. Inst.*, **I**, 70 pp.
- WILSON, B. R. & K. GILLETT 1971: *Australian Shells* – A. H. & A. W. Reed, Sydney. 168 pp., 106 coloured plates.
- YONGE, C. M. 1953: Observations on *Hipponix antiquatus* (Linnaeus). – *Proc. California Acad. Sci.*, **XXVIII**(1): 1–24, 9 text figs.

Dr. Al Barash,
Dept. of Zoology,
Tel Aviv University
Ramat-Aviv
69978 Tel-Aviv, Israel